## Late-Stage Colorectal Cancer Incidence, by Year, Maine, 2004-2009

Year	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
2004	421	32.2	29.2 - 35.4	27.3	24.8 - 30.1
2005	402	30.6	27.7 - 33.8	26.0	23.5 - 28.6
2006	383	29.1	26.3 - 32.2	24.0	21.6 - 26.5
2007	409	31.0	28.1 - 34.2	25.3	22.9 - 27.9
2008	371	28.1	25.3 - 31.1	22.7	20.4 - 25.1
2009	388	29.4	26.6 - 32.5	23.5	21.2 - 26.0

## Late-Stage Colorectal Cancer Incidence, by 3-year averages, Maine, 2004-2009

Year	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
2004-2006	1,206	30.6	28.9 - 32.4	25.7	24.3 - 27.3
2005-2007	1,194	30.3	28.6 - 32.0	25.1	23.7 - 26.6
2006-2008	1,163	29.4	27.8 - 31.2	24.0	22.6 - 25.4
2007-2009	1,171	29.6	27.9 - 31.4	23.9	22.5 - 25.3

Data Source: Maine Cancer Registry, November 2011 NPCR data submission.

Count: Number of cases diagnosed with malignant cancer during that time period.

Late-Stage: Cancers with 'regional' or 'distant' spread according to 2000 Derived Summary Stage criteria.

Colorectal Cancer: SEER cancer site recodes: 21041-21049,21051,21052 ( ICD-O-3 codes C180-C189,C260,C199,C209).

Crude rates are new cases per 100,000 population.

95% CI: 95% confidence interval of the rate.

Rates are calculated using SEER\*Stat Version 7.0.5.